



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the equation to a cone having (x', y', z') for vertex, and (3) for base.

$$(8) \text{ is } \frac{z'^2}{a^2}x^2 + \frac{z'^2}{b^2}y^2 + \left(\frac{x'^2}{a^2} + \frac{y'^2}{b^2} - 1\right)z^2 - \frac{2y'z'}{b^2}yz - \frac{2x'z'}{a^2}xz + 2z'z - z'^2 = 0 \dots (9).$$

The conditions for (9) to be a cone of revolution are $x'=0$,

$$\frac{z'^2}{a^2} - \frac{y'^2}{b^2 - a^2} = 1 \dots (10),$$

an hyperbola for the required locus.

Also solved by G. B. M. ZERR.

100. Proposed by CHARLES CARROLL CROSS, Libertytown, Md.

O, O_1, O_2, O_3 are the centers of the inscribed and three escribed circles of a triangle ABC . Prove $AO \cdot AO_1 \cdot AO_2 \cdot AO_3 = AB^2 \cdot AC^2$.

I. Solution by the PROPOSER.

Consider the ex-central triangle ABC as the original triangle. Then $H_a H_b H_c$ is the pedal triangle, and the incenter O becomes the orthocenter H .

Hence we have to prove $AH_c \times BH_c \times CH_c \times HH_c = H_a H_c^2 \times H_b H_c^2$.

We readily find by trigonometry that

$$AH_c = \frac{b^2 + c^2 - a^2}{2c}, \quad BH_c = \frac{a^2 + c^2 - b^2}{2c},$$

$$CH_c = \frac{\sqrt{[4a^2c^2 - (a^2 + c^2 - b^2)^2]}}{2c} = \frac{2\Delta}{c}, \quad HH_c = \frac{(b^2 + c^2 - a^2)(a^2 + c^2 - b^2)}{8c\Delta},$$


$$H_a H_c = \frac{b(a^2 + c^2 - b^2)^2}{2ac}, \quad H_b H_c = \frac{a(b^2 + c^2 - a^2)}{2bc}.$$

Substituting in the problem we have

$$\begin{aligned} \frac{b^2 + c^2 - a^2}{2c} \times \frac{a^2 + c^2 - b^2}{2c} \times \frac{2\Delta}{c} \times \frac{(b^2 + c^2 - a^2)(a^2 + c^2 - b^2)}{8c\Delta} \\ = \frac{b^2(a^2 + c^2 - b^2)^2}{4a^2c^2} \times \frac{a^2(b^2 + c^2 - a^2)^2}{4b^2c^2}. \end{aligned}$$


Since these equations cancel, the proposition is proved.

Mr. Cross should have been credited with solutions of problems 96 and 98 in Geometry, and 100 and 101 in Arithmetic.



II. Solution by WALTER H. DRANE, Graduate Student at Harvard University, and J. SCHEFFER, A. M., Hagerstown, Md.

Let ABC be the given triangle, O, O_1, O_2, O_3 , the centers of the inscribed and the three escribed circles. Then $O_1BO_2, O_1AO_3, O_2CO_3$, and AOO_2 are straight lines; also OB, OA, OC are each perpendicular to O_1BO_2, O_1AO_3 , and O_2CO_3 , respectively. In triangles AOC and BAO_2 , $\angle OAC = \angle BAO_2$ and $\angle OCA = \angle BO_2A$ since we have $\angle OCA = 90^\circ - \angle ACO_3 = 90^\circ - \frac{1}{2}(\angle CAB + \angle CBA) = 90^\circ - (\angle OAB + \angle OBA) = 90^\circ - [180^\circ - (\angle O_1AB + \angle O_1BA)] = 90^\circ - \angle O = \angle O_1O_2A$.



\therefore triangles AOC and ABO_2 are similar, and we have

$$AO : AB :: AC : AO_2 \dots \dots \dots (1).$$

Again in triangles O_1BA and AO_3C , $\angle O = \angle ACO_3$ and $\angle O_3AC = \angle O_1AB$. Hence triangles O_1BA and AO_3C are similar, and we have,

$$AO_1 : AC :: AB : AO_3 \dots \dots \dots (2).$$

Multiplying (1) by (2) $AO.AO_1 : AB.AC :: AB.AC : AO_2.AO_3$.

$$\therefore AO.AO_1.AO_2.AO_3 = AB^2.AC^2.$$

Q. E. D.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

In the figure of the last solution, draw the lines OD and O_1D_1 perpendicular to AC . Then $AO = \sqrt{(OD^2 + AD^2)} = \sqrt{(r^2 + r^2 \cot^2 \frac{1}{2}A)} = r \operatorname{cosec} \frac{1}{2}A$.

Similarly, $AO_1 = r_1 \operatorname{cosec} \frac{1}{2}A$.

$$AO_2 = \sqrt{(OO_2^2 - AO^2)} = AO \sqrt{[(OO_2^2/AO^2) - 1]} = AO \cot \frac{1}{2}C.$$

Similarly, $AO_3 = AO \cot \frac{1}{2}B$.


$$\begin{aligned} \therefore AO.AO_1.AO_2.AO_3 &= r^3 r_1 \operatorname{cosec}^4 \frac{1}{2}A \cot \frac{1}{2}B \cot \frac{1}{2}C \\ &= (s-a)(s-b)(s-c) \operatorname{cosec}^4 \frac{1}{2}A \tan^2 \frac{1}{2}A \\ &= s(s-a)(s-b)(s-c) / \sin^2 \frac{1}{2}A \cos^2 \frac{1}{2}A \\ &= 4s(s-a)(s-b)(s-c) / \sin^2 A = 4S^2 / \sin^2 A. \end{aligned}$$

$$\text{But } \sin A = 2S/bc = 2S/AB.AC. \therefore AO.AO_1.AO_2.AO_3 = AB^2.AC^2.$$

Also solved by ELMER SCHUYLER.

101. Proposed by E. W. MORRELL, A. M., Late Professor of Mathematics, Montpelier Seminary, Montpelier, Vt.

AB is the diameter of a circle and Q_0 any point on the circumference; $Q_1, Q_2, Q_3 \dots$ are the points of bisection of the arcs $AQ_0, AQ_1, AQ_2 \dots$. Prove that $BQ_1, BQ_2, BQ_3 \dots BQ_n = OA^n.(AQ_0/AQ_n)$.



Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.; J. SCHEFFER, A. M., Hagerstown, Md., and ELMER SCHUYLER, High Bridge, N. J.

Let O be the center of the circle.

$$\angle ABQ_0 = \theta.$$

$$\therefore BQ_1 = AB \cos \frac{1}{2}\theta, BQ_2 = AB \cos(\theta/2^2).$$

$$BQ_3 = AB \cos(\theta/2^3), BQ_n = AB \cos(\theta/2^n).$$